

# PATENT ABSTRACTS OF JAPAN

(11)Publication number : 2000-290262

(43)Date of publication of application : 17.10.2000

(51)Int.Cl.

C07D239/94

(21)Application number : 2000-091300

(71)Applicant : PFIZER PROD INC

(22)Date of filing : 29.03.2000

(72)Inventor : LEHNER RICHARD SHELTON  
NORRIS TIMOTHY  
SANTAFIANOS DINOS PAUL

(30)Priority

Priority number : 99 127072

Priority date : 31.03.1999


Priority country : US

## (54) METHOD AND INTERMEDIATE FOR PRODUCING ANTICANCER COMPOUND


(57)Abstract:

PROBLEM TO BE SOLVED: To obtain an anticancer compound in a high yield useful for treating a highly proliferative disorder such as cancer in mammals by using a specific synthetic intermediate.

SOLUTION: (A) A compound of formula I [R1 and R2 are each a 1-10C alkyl, a 1-10C alkoxy, preferably R1 and R2 are each 2-methoxyethoxy; R15 is H, a 1-10C alkyl or the like, preferably H; G is C(OH)R3R4 or SiR3R4R5 (R3 to R5 are each a 1-6C alkyl)] is treated, (B) in the case of G being C(OH)R3 R4, with an alkaline (earth) metal hydroxide in a hydroxy-substituted 1-10C alkyl-containing solvent or, (C) in the case of G being SiR3R4R5, with a tetra (1-6C alkyl) ammonium fluoride compound in an aprotic solvent to give a compound of formula II. The compound of formula I can be obtained by treating a compound of formula III with a compound of formula IV.




I



II



III



IV

(19) 日本国特許庁 (J P)

(12) 特 許 公 報 (B 2)

(11) 特許番号

特許第3420549号

(P3420549)

(45) 発行日 平成15年6月23日 (2003. 6. 23)

(24) 登録日 平成15年4月18日 (2003. 4. 18)

(51) Int.Cl.<sup>7</sup>

識別記号

F I

C 0 7 D 239/94

C 0 7 D 239/94

請求項の数14(全 14 頁)

(21) 出願番号 特願2000-91300(P2000-91300)

(22) 出願日 平成12年3月29日 (2000. 3. 29)

(65) 公開番号 特開2000-290262(P2000-290262A)

(43) 公開日 平成12年10月17日 (2000. 10. 17)

審査請求日 平成12年3月29日 (2000. 3. 29)

(31) 優先権主張番号 6 0 / 1 2 7 0 7 2

(32) 優先日 平成11年3月31日 (1999. 3. 31)

(33) 優先権主張国 米国 (U S)

前置審査

(73) 特許権者 397067152

ファイザー・プロダクツ・インク

アメリカ合衆国コネチカット州グロトン

市イースタン・ポイント・ロード

(72) 発明者 リチャード・シェルトン・ラーナー

アメリカ合衆国コネチカット州06339,

レッドヤード, ウィンドワード・レイン

12

(72) 発明者 ティモシー・ノリス

アメリカ合衆国コネチカット州06335,

ゲールズ・フェリー, フライアー・タッ

ク・ドライブ 27

(74) 代理人 100089705

弁理士 社本 一夫 (外5名)

審査官 内藤 伸一

最終頁に続く


(54) 【発明の名称】 抗癌性化合物を製造するための方法と中間体

1

2

.....  
.....  
.....

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....




1


10

.....  
.....

(2)



4





10


\*20\*

\*30\*

\*40\*





6




$R^1$


$R^{15}$

$G$







48




49




50




51




52




53




54





55








30



40




50



30

40

50



30

40

50





\*10\*

\*20\*





\*30\*

\*40\*

50







20

30

40

50

[illegible]

$\delta_H$  (300 MHz;  $CDCl_3$ ) 0.24 (9H, s), 3.56 (2H, bs), 6.82 (1H, ddd,  $J = 1.0, 2.3$  & 8.0), 6.76 (1H, t,  $J = 2.2$ ), 6.87 (1H, dt,  $J = 7.7$  & 1.2), 7.07 (1H, t,  $J = 7.5$ );  $\delta_C$  (75.5 MHz;  $CDCl_3$ ) 93.4, 105.4, 115.6, 118.2, 122.4, 123.8, 129.2, 146.2;  $m/e$  190 ( $M+H$ )<sup>+</sup>

10

$\delta_H$  (400 MHz;  $CDCl_3$ ) 0.21 (9H, s), 3.38 (5H, s), 3.41 (3H, s), 3.72 (2H, m), 3.77 (2H, m), 4.10 (2H, s), 4.53 (2H, s), 7.20 (1H, t,  $J = 7.8$ ), 7.23-7.28 (2H, m), 7.75 (1H, d,  $J = 7.8$ ), 7.88 (1H, s), 8.20 (1H, s), 8.42 (1H, s);  $m/e$  466 ( $M+H$ )<sup>+</sup>

20

$\delta_H$  (300 MHz;  $d_6$ -DMSO) 3.39 (6H, s), 3.77-3.80 (4H, m), 4.30 (1H, s), 7.39 (1H, s), 7.41 (1H, d,  $J = 7.9$ ), 7.60 (1H, t,  $J = 7.8$ ), 7.79 (1H, d,  $J = 8.1$ ), 7.88 (1H, s), 8.40 (1H, s), 8.86 (1H, s), 11.48 (1H, bs);  $\delta_C$  (100 MHz;  $d_6$ -DMSO) 58.4, 58.5, 69.7, 69.2, 69.7, 57.6, 81.3, 83.0, 100.3, 105.2, 107.2, 121.9, 125.4, 127.6, 128.9, 129.2, 135.2, 137.7, 146.3, 149.2, 155.4, 166.0;  $m/e$  394 ( $M+H$ )<sup>+</sup>

40

$\delta_H$  (400 MHz;  $d_6$ -DMSO) 1.44 (6H, s), 3.31-3.32 (5H, m), 3.65-3.75 (2H, m), 4.24-4.30 (2H, m), 4.35-4.37 (2H, m), 7.25 (1H, m), 7.39 (2H, m), 7.72-7.74 (2H, m), 8.47 (1H, s), 8.79 (1H, s), 11.64 (1H, s);  $m/e$  452 ( $M+H$ )<sup>+</sup>

50

.....

.....

.....

.....

[illegible]



$\delta_H$  (300 MHz;  $CDCl_3$ ) 1.17 (3H, t, J 7.6), 2.56 (2H, q, J 7.6), 3.33 (3H, s), 3.55-3.66 (2H, m), 4.07-4.11 (2H, m), 5.11 (2H, s), 6.83 (1H, d, J 7.7), 7.18-7.29 (5H, m), 7.35-7.42 (4H, m), 7.50 (1H, d, J 8.0), 8.20 (1H, bs), 8.51 (1H, s);  $\delta_C$  (75.5 MHz;  $CDCl_3$ ) 14.2, 15.4, 28.8, 59.2, 69.2, 70.7, 70.8, 103.2, 106.1, 108.4, 119.7, 121.7, 124.0, 127.3, 128.1, 128.5, 128.8, 135.8, 138.6, 145.1, 147.0, 148.5, 153.7, 154.2, 156.9;  $\nu_{max}$  (KBr)  $cm^{-1}$  1625, 1611, 1576;  $m/z$  430 (M+H)<sup>+</sup>; { 实验值: C, 71.42; H, 6.50; N, 9.48.  $C_{26}H_{27}N_3O_3$  计算值: C, 72.70; H, 6.34; N, 9.78%}.

$\delta_H$  (300 MHz;  $CDCl_3$ ) 0.93 (3H, t, J 7.4), 1.19 (3H, t, J 7.6), 1.45 (2H, 次亚峰 J 7.5), 1.76 (2H, 王峰, J 6.9), 2.61 (2H, q, J 7.6), 3.38 (3H, s), 3.70-3.74 (2H, m), 4.00 (2H, t, J 6.6), 4.12-4.16 (2H, m), 6.94 (1H, d, J 7.7), 7.16 (1H, s), 7.24 (1H, t, J 7.5), 7.34 (1H, s), 7.44 (1H, s), 7.51 (1H, d, J 8.0), 7.85 (1H, bs), 8.60 (1H, s);  $\delta_C$  (75.5 MHz;  $CDCl_3$ ) 13.8, 15.4, 18.2, 28.8, 30.8, 59.3, 69.7, 69.9, 70.9, 103.2, 108.2, 108.9, 119.6, 121.6, 124.6, 128.9, 138.6, 145.2, 147.2, 149.8, 153.6, 154.9, 156.8;  $\nu_{max}$  (KBr)  $cm^{-1}$  1618, 1576, 1519;  $m/z$  395 (M+H)<sup>+</sup>; { 实验值: C, 70.60; H, 7.56; N, 10.66.  $C_{22}H_{25}N_3O_3$  计算值: C, 69.85; H, 7.39; N, 10.63%}.

$\delta_H$  (300 MHz;  $CDCl_3$ ) 3.31 (3H, s), 3.35 (3H, s), 3.62-3.65 (2H, m), 3.70-3.72 (2H, m), 3.74 (5H, s), 4.04-4.11 (4H, m), 6.83 (2H, d, J 9.0), 7.09 (1H, s), 7.33 (1H, s), 7.46 (2H, d, J 9.0), 8.12 (1H, bs), 8.1H, s);  $\delta_C$  (75.5 MHz;  $CDCl_3$ ) 55.4, 59.2, 68.2, 69.3, 70.4, 70.8, 103.1, 108.3, 109.1, 114.2, 124.7, 131.4, 146.3, 148.9, 153.7, 154.3, 156.7, 157.3;  $\nu_{max}$  (KBr)  $cm^{-1}$  1619, 1600, 1562, 1511;  $m/z$  400 (M+H)<sup>+</sup>; { 实验值: C, 63.38; H, 6.37; N, 10.47.  $C_{21}H_{23}N_3O_3$  计算值: C, 63.42; H, 6.31; N, 10.52%}.

$\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 3.34 (3H, s), 3.91 (2H, t, J 4.2), 3.74 (3H, s), 4.10 (2H, bs), 5.13 (2H, s), 5.83 (2H, c, J 8.9), 7.20-7.30 (5H, m), 7.36-7.38 (3H, m), 7.47 (2H, d, J 8.5), 8.10 (1H, bs), 8.54 (1H, s);  $\delta_{\text{C}}$  (75.5 MHz;  $\text{CDCl}_3$ ) 55.5, 59.3, 60.2, 70.7, 70.3, 103.3, 109.0, 109.1, 114.2, 124.8, 127.3, 128.1, 128.5, 131.3, 135.8, 146.9, 149.6, 153.7, 154.2, 154.2, 156.6, 157.2;  $\nu_{\text{max}}$  (KBr)  $\text{cm}^{-1}$  1619, 1580, 1511;  $m/z$  432 (M+H) $^{+}$ ; (元素分析 C, 69.46; H, 5.85; N, 9.68).  $\text{C}_{22}\text{H}_{25}\text{N}_3\text{O}_4$  計算値: C, 69.59; H, 5.84; N, 9.74%.

$\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 3.33 (3H, s), 3.39 (3H, s), 3.42-3.45 (2H, m), 3.48-3.51 (2H, m), 3.59 (3H, s), 3.74-3.78 (2H, m), 4.18-4.20 (2H, m), 8.38 (1H, s), 7.11-7.20 (4H, m), 7.83 (2H, t, J 7.8), 8.66 (1H, s);  $\delta_{\text{C}}$  (75.5 MHz;  $\text{CDCl}_3$ ) 42.0, 55.2, 59.3, 67.6, 68.2, 70.3, 70.4, 106.6, 107.9, 110.9, 125.8, 126.0, 126.9, 147.0, 148.4, 148.7, 153.0, 153.4, 160.4;  $\nu_{\text{max}}$  (KBr)  $\text{cm}^{-1}$  1615, 1671, 1497;  $m/z$  384 (M+H) $^{+}$ ; (元素分析 C, 55.65; H, 5.52; N, 11.01).  $\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_4$  計算値: C, 55.78; H, 5.57; N, 10.96%.

$\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 0.93 (3H, t, J 7.4), 1.45 (2H, 六重線 J 7.4), 1.80 (2H, 五重線, J 6.7), 3.25 (3H, s), 3.44-3.52 (4H, m), 3.59 (3H, s), 4.05 (2H, t, J 6.7), 6.34 (1H, s), 7.12-7.21 (4H, m), 7.34 (2H, t, J 7.7), 8.69 (1H, s);  $\delta_{\text{C}}$  (75.5 MHz;  $\text{CDCl}_3$ ) 13.8, 19.2, 30.7, 42.0, 59.2, 67.6, 68.6, 70.4, 106.5, 107.7, 110.6, 125.8, 125.9, 129.9, 147.0, 148.6, 153.0, 153.6, 160.4;  $\nu_{\text{max}}$  (KBr)  $\text{cm}^{-1}$  1616, 1572, 1543;  $m/z$  382 (M+H) $^{+}$ ; (元素分析 C, 69.39; H, 7.38; N, 10.86).  $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_5$  計算値: C, 69.27; H, 7.14; N, 11.02%.